

Otterbein University

Digital Commons @ Otterbein

Chemistry Faculty Scholarship

Chemistry

3-1-2012

Tris(3-nitropentane-2,4-dionato-κ O,O')-cobalt(III)

Dean Johnston

Follow this and additional works at: https://digitalcommons.otterbein.edu/chem_fac

 Part of the [Chemistry Commons](#)

Repository Citation

Johnston, Dean, "Tris(3-nitropentane-2,4-dionato-κ O,O')-cobalt(III)" (2012). *Chemistry Faculty Scholarship*. 9.

https://digitalcommons.otterbein.edu/chem_fac/9

This Article is brought to you for free and open access by the Chemistry at Digital Commons @ Otterbein. It has been accepted for inclusion in Chemistry Faculty Scholarship by an authorized administrator of Digital Commons @ Otterbein. For more information, please contact digitalcommons07@otterbein.edu.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/221698220>

Tris(3-nitropentane-2,4-dionato- κ O,O')-cobalt(III)

Article in *Acta Crystallographica Section E Structure Reports Online* · March 2012

Impact Factor: 0.35 · DOI: 10.1107/S160053681200668X · Source: PubMed

CITATION

1

READS

379

3 authors, including:



Dean H Johnston

Otterbein University

29 PUBLICATIONS 908 CITATIONS

SEE PROFILE

Tris(3-nitropentane-2,4-dionato- κ^2O,O')-cobalt(III)

Dean H. Johnston,* Jack T. Brangham and Christopher D. Rapp

Department of Chemistry, Otterbein University, Westerville, OH 43081, USA

Correspondence e-mail: djohnston@otterbein.edu


Received 4 February 2012; accepted 14 February 2012

Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(C-C) = 0.004$ Å; disorder in main residue; R factor = 0.026; wR factor = 0.061; data-to-parameter ratio = 10.3.

The structure of the title compound, $[Co(C_5H_6NO_4)_3]$, consists of a Co^{III} ion octahedrally coordinated by three bidentate 3-nitropentane-2,4-dionate ligands. The complex was prepared *via* the nitration of tris(2,4-pentanedionato- κ^2O,O')cobalt(III) with a solution of copper(II) nitrate in glacial acetic acid. The central C atom and the nitro group of one 3-nitropentane-2,4-dionate ligand are disordered over two positions with an occupancy ratio of 0.848 (4):0.152 (4). A second nitro group is also disordered over two orientations with an occupancy ratio of 0.892 (7):0.108 (7). Two of the ligand methyl groups form $C-H \cdots O$ interactions with two different nitro groups to form chains running along the c axis. Additional $C-H \cdots O$ interactions are found between ligand methyl groups and the cobalt-bound O atoms, also resulting in the formation of chains along the c axis.

Related literature

For the preparation of derivatized tris(2,4-pentanedionato) metal complexes, see: Collman *et al.* (1962, 1963); Collman (1965); Schirado *et al.* (1971); James (1974); Shalhoub (1980). For spectroscopic properties of the title compound, see: Singh & Sahai (1967, 1968); Larsson & Eskilsson (1969); Fleming & Thorton (1973, 1975); Tsiamis *et al.* (1987). For crystallographic studies of related compounds, see: Appleton *et al.* (1992); Abrahams *et al.* (1998); Tsiamis *et al.* (1998); von Chrzanowski *et al.* (2007). For a review of graph-set analysis of hydrogen-bonding patterns, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$[Co(C_5H_6NO_4)_3]$
 $M_r = 491.25$
 Tetragonal, $I4_1cd$
 $a = 32.7078$ (18) Å
 $c = 7.4976$ (6) Å
 $V = 8020.9$ (9) Å³

$Z = 16$
 Mo $K\alpha$ radiation
 $\mu = 0.93$ mm⁻¹
 $T = 200$ K
 $0.48 \times 0.40 \times 0.32$ mm

Data collection

Bruker SMART X2S benchtop diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{min} = 0.665$, $T_{max} = 0.756$

24724 measured reflections
 3393 independent reflections
 3151 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$
 $wR(F^2) = 0.061$
 $S = 1.04$
 3393 reflections
 330 parameters
 159 restraints

H-atom parameters constrained
 $\Delta\rho_{max} = 0.20$ e Å⁻³
 $\Delta\rho_{min} = -0.19$ e Å⁻³
 Absolute structure: Flack (1983),
 1466 Friedel pairs
 Flack parameter: 0.003 (12)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$C1-H1B \cdots O7^i$	0.98	2.33	3.087 (4)	134
$C11-H11B \cdots O12^{ii}$	0.98	2.55	3.240 (4)	128
$C10-H10C \cdots O5^{iii}$	0.98	2.46	3.433 (3)	176
$C15-H15C \cdots O3^{iv}$	0.98	2.57	3.542 (4)	174

Symmetry codes: (i) $-x + 1, y, z - \frac{1}{2}$; (ii) $x, -y + 1, z - \frac{1}{2}$; (iii) $-y + 1, x - \frac{1}{2}, z + \frac{1}{4}$; (iv) $x, y, z + 1$.

Data collection: *GIS* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) and *OLEX2* (Dolomanov *et al.*, 2009); molecular graphics: *PLATON* (Spek, 2009), *Mercury* (Macrae *et al.*, 2008) and *POV-RAY* (Cason, 2004); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

This work was supported in part by the National Science Foundation through grant No. CHE-0942850.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZL2451).

References

- Abrahams, B. F., Hoskins, B. F., McFadyen, D. W. & Perrin, L. C. (1998). *Acta Cryst.* **C54**, 1807–1809.
- Appleton, T. G., Gahan, L. R. & Oliver, P. J. (1992). *Aust. J. Chem.* **45**, 797–805.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N. L. (1995). *Angew. Chem. Int. Ed.* **34**, 1555–1573.
- Bruker (2009). *GIS, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cason, C. J. (2004). *POV-RAY*. Persistence of Vision Raytracer Pty Ltd, Williamstown, Victoria, Australia.
- Chrzanowski, L. S. von, Lutz, M. & Spek, A. L. (2007). *Acta Cryst.* **C63**, m283–m288.
- Collman, J. P. (1965). *Angew. Chem. Int. Ed.* **4**, 132–138.
- Collman, J. P., Goldby, S., Young, W. L. III & Marshall, R. (1962). *Inorg. Chem.* **1**, 704–710.
- Collman, J. P., Young, W. L. III & Kauffman, G. B. (1963). *Inorg. Synth.* **7**, 205–207.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Fleming, C. A. & Thorton, D. A. (1973). *J. Mol. Struct.* **17**, 79–89.
- Fleming, C. A. & Thorton, D. A. (1975). *J. Mol. Struct.* **25**, 271–279.
- James, B. D. (1974). *J. Chem. Educ.* **51**, 568.
- Larsson, R. & Eskilsson, O. (1969). *Acta Chem. Scand.* **23**, 1765–1779.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Schirado, T., Gennari, E., Merello, R., Decinti, A. & Bunel, S. (1971). *J. Inorg. Nucl. Chem.* **33**, 3417–3426.
- Shalhoub, G. M. (1980). *J. Chem. Educ.* **57**, 525–528.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Singh, P. R. & Sahai, R. (1967). *Aust. J. Chem.* **20**, 649–655.
- Singh, P. R. & Sahai, R. (1968). *Inorg. Nucl. Chem. Lett.* **4**, 513–516.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Tsiamis, C., Cambanis, S. & Hadjikostas, C. (1987). *Inorg. Chem.* **26**, 26–32.
- Tsiamis, C., Stergiou, A. C., Anesti, V., Blaton, N. M. & Peeters, O. M. (1998). *Inorg. Chim. Acta*, **269**, 332–336.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, m312–m313 [doi:10.1107/S160053681200668X]

Tris(3-nitropentane-2,4-dionato- κ^2O,O')cobalt(III)**Dean H. Johnston, Jack T. Brangham and Christopher D. Rapp****Comment**

The electrophilic substitution chemistry of the 2,4-pentanedionato (acetylacetonate, or acac) ligand has been studied for many years (Collman, *et al.*, 1962; Collman, *et al.*, 1963; Collman, 1965; Schirado, *et al.*, 1971), but relatively few of these derivatives have been studied crystallographically, especially for the tri-substituted complexes. The nitro derivative of the cobalt complex is readily prepared and its synthesis and characterization have been described as part of several educational laboratory activities (James, 1974; Shalhoub, 1980).

The average cobalt-oxygen bond length in the title compound is 1.869 (4) Å, slightly shorter than the average cobalt-oxygen bond length observed for the [Co(acac)₃] complex determined at a similar temperature (von Chrzanowski, *et al.*, 2007). All three nitro groups are twisted with respect to their 2,4-pentanedionato ligands (Fig. 1), forming angles of 49.3 (1), 59.3 (2), and 50.3 (2) degrees for the major components and 67.2 (2) and 51.6 (8) degrees for the minor disorder components. These are similar to the angle of 50.7 degrees observed for the mono-nitro cobalt complex (Appleton *et al.*, 1992). The disorder in the positioning of one chelate ring has been observed previously (as large thermal parameters) for analogous complexes of cobalt and manganese (Appleton *et al.*, 1992).

Analysis of packing (Fig. 2) and close contacts shows two different types of C—H \cdots O interactions (Table 1). The first type, shown in Figure 3(a) and 3(b), forms between methyl group hydrogen atoms and the nitro group on an adjacent molecule. The second type of C—H \cdots O, shown in Figure 3(c) and 3(d), forms between methyl group hydrogen atoms and the cobalt-bound oxygen atom on an adjacent molecule. This second type of interaction is commonly seen in 2,4-pentanedionato complexes (von Chrzanowski *et al.*, 2007). These hydrogen-bonding interactions result in the formation of four different types of C(6) chains (Bernstein *et al.*, 1995), shown in Figure 4(a) through 4(d). In all four cases, the primary direction of the chain is along the *c* axis.

Experimental

The complex was prepared according to the procedure of Collman *et al.* (1963). Approximately 5.37 g (0.023 mol) of finely ground copper(II) nitrate trihydrate was mixed with 100 ml (1.06 mol) of acetic anhydride. Cobalt(III) acetylacetonate (2.5 g, 0.0070 mol) was added to the mixture and stirred with cooling for approximately two hours. A combination of water (300 ml), ice (300 g), and sodium acetate (7.5 g, 0.055 mol) was then added and the mixture was stirred for an additional two hours. The dark-green precipitate was vacuum filtered and washed with water and cold ethanol. The crude product was recrystallized from boiling chloroform and hot ethanol. The final product consisted of large, dark green crystals that were obtained in an overall yield of 62% (2.14 g).

The IR spectrum (ATR cell) displayed strong peaks at 1561 cm⁻¹ (ν_{ring}), 1518 cm⁻¹ (ν_{as} , NO₂), 1341 cm⁻¹ (ν_{s} , NO₂), and 825 cm⁻¹ ($\delta_{\text{C-H}}$). Raman spectra (532 nm excitation) gave strong peaks at 1345 cm⁻¹ (ν_{s} , NO₂), 828 cm⁻¹ ($\delta_{\text{C-H}}$), 470 cm⁻¹ and 450 cm⁻¹ ($\nu_{\text{Co-O}}$).

Refinement

All hydrogen atoms were located in the difference map and refined with the atom positions constrained to an ideal tetrahedron with C—H distances of 0.98 Å. A riding model was used for all hydrogen atoms with $U_{\text{iso}}(\text{H}) = 1.5$ times $U_{\text{iso}}(\text{C})$.

One of the 3-nitropentane-2,4-dionato ligands was modeled as disordered over two positions for four atoms, C13/C13A, N3/N3A, O11/O11A, and O12/O12A and refined to give an occupancy ratio of 0.848 (4):0.152 (4). Carbon-carbon distances between similar atoms in the disordered ligand were restrained to be similar within a standard deviation of 0.02 Å. The nitro groups and their respective carbon atoms (C13/N3/O11/O12, C13A/N3A/O11A/O12A) were restrained to lie in a common plane, as were atoms C12, C13A, C14 and N3A. The anisotropic displacement parameters for the atom pairs N3/N3A and C13/C13A were constrained to be the same. The nitro group on a second ligand (N2, O9, O10) was modeled as a disordered group over two orientations and refined to give an occupancy ratio of 0.892 (7):0.108 (7).

Anisotropic displacement parameters were restrained to be similar (standard deviations of 0.01 Å², 0.02 Å²) for 1,2 and 1,3-bonded atoms and approximately isotropic (standard deviation of 0.1 Å²) for all disordered oxygen atoms. Anisotropic displacement parameters were also restrained to be similar (with a standard deviation of 0.01 Å²) for all atoms within the disordered nitro groups.

Computing details

Data collection: GIS (Bruker, 2009); cell refinement: *SAINTE* (Bruker, 2009); data reduction: *SAINTE* (Bruker, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) and *OLEX2* (Dolomanov *et al.*, 2009); molecular graphics: *PLATON* (Spek, 2009), Mercury (Macrae *et al.*, 2008) and *POV-RAY* (Cason, 2004); software used to prepare material for publication: *publCIF* (Westrip, 2010).






Figure 1


The molecular structure of the title compound showing the atom labeling scheme and drawn with 50% probability displacement ellipsoids for non-H atoms.

**Figure 2**

The packing of the title compound viewed along the c axis. The minor components of the disordered ligands are not shown.

**Figure 3**

Hydrogen-bond interactions in the title compound. The C—H...O contacts are shown with dashed lines. (a) The C1—H1B...O7ⁱ contact, (b) the C11—H11B...O12ⁱⁱ contact, (c) the C10—H10C...O5ⁱⁱⁱ contact, (d) the C15—H15C...O3^{iv} contact. [Symmetry codes: (i) $-x + 1, y, z - 1/2$; (ii) $x, -y + 1, z - 1/2$; (iii) $-y + 1, x - 1/2, z + 1/4$; (iv) $x, y, z + 1$.]


Figure 4

The four different types of $C(6)$ hydrogen-bonded chains formed by the title compound. The $C—H\cdots O$ contacts are shown with dashed lines. (a) The $C1—H1B\cdots O7^i$ chain, viewed along the b axis, (b) the $C11—H11B\cdots O12^{ii}$ chain, viewed along the a axis, (c) the $C10—H10C\cdots O5^{iii}$ chain, viewed along the c axis (d) the $C15—H15C\cdots O3^{iv}$ chain, viewed along the a axis. [Symmetry codes: (i) $-x + 1, y, z - 1/2$; (ii) $x, -y + 1, z - 1/2$; (iii) $-y + 1, x - 1/2, z + 1/4$; (iv) $x, y, z + 1$.]

Tris(3-nitropentane-2,4-dionato- $\kappa^2 O, O'$)cobalt(III)

Crystal data

$[\text{Co}(\text{C}_5\text{H}_6\text{NO}_4)_3]$
 $M_r = 491.25$
 Tetragonal, $I4_1cd$
 $a = 32.7078$ (18) Å
 $c = 7.4976$ (6) Å
 $V = 8020.9$ (9) Å³
 $Z = 16$
 $F(000) = 4032$

$D_x = 1.627$ Mg m⁻³
 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 8816 reflections
 $\theta = 2.5$ – 24.6°
 $\mu = 0.93$ mm⁻¹
 $T = 200$ K
 Block, green
 $0.48 \times 0.40 \times 0.32$ mm

Data collection

Bruker SMART X2S benchtop
 diffractometer
 Radiation source: fine-focus sealed tube
 Doubly curved silicon crystal monochromator
 Detector resolution: 8.3330 pixels mm⁻¹
 φ and ω scans
 Absorption correction: multi-scan
 (*SADABS*; Bruker, 2009)
 $T_{\min} = 0.665$, $T_{\max} = 0.756$

24724 measured reflections
 3393 independent reflections
 3151 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$
 $\theta_{\max} = 25.1^\circ$, $\theta_{\min} = 2.5^\circ$
 $h = -38 \rightarrow 38$
 $k = -38 \rightarrow 34$
 $l = -7 \rightarrow 8$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.026$

$wR(F^2) = 0.061$
 $S = 1.04$
 3393 reflections

330 parameters
 159 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map
 Hydrogen site location: difference Fourier map
 H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0316P)^2 + 2.8577P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.20 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.19 \text{ e } \text{Å}^{-3}$
 Absolute structure: Flack (1983), 1466 Friedel
 pairs
 Flack parameter: 0.003 (12)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Co1	0.623604 (9)	0.363125 (9)	0.26098 (5)	0.02596 (9)	
O1	0.60216 (5)	0.41367 (5)	0.1946 (3)	0.0344 (4)	
O2	0.57311 (5)	0.33960 (5)	0.3185 (2)	0.0328 (4)	
C1	0.55984 (10)	0.46990 (8)	0.1698 (5)	0.0506 (8)	
H1A	0.5562	0.4848	0.2820	0.076*	
H1B	0.5357	0.4737	0.0944	0.076*	
H1C	0.5841	0.4803	0.1076	0.076*	
C2	0.56542 (8)	0.42488 (7)	0.2088 (4)	0.0335 (6)	
C3	0.53378 (7)	0.39769 (8)	0.2526 (4)	0.0343 (6)	
C4	0.53853 (8)	0.35653 (8)	0.3056 (3)	0.0321 (6)	
C5	0.50367 (9)	0.32949 (9)	0.3579 (4)	0.0457 (7)	
H5A	0.4921	0.3167	0.2511	0.069*	
H5B	0.4826	0.3458	0.4174	0.069*	
H5C	0.5135	0.3082	0.4394	0.069*	
N1	0.49174 (8)	0.41312 (8)	0.2502 (4)	0.0537 (7)	
O7	0.48554 (8)	0.44588 (8)	0.3244 (4)	0.0868 (9)	
O8	0.46521 (7)	0.39291 (9)	0.1762 (4)	0.0736 (8)	
O3	0.61607 (5)	0.34505 (5)	0.0263 (3)	0.0323 (4)	
O4	0.64932 (5)	0.31545 (5)	0.3405 (2)	0.0292 (4)	
C6	0.61730 (9)	0.30447 (8)	-0.2281 (4)	0.0458 (7)	
H6A	0.5967	0.3245	-0.2651	0.069*	
H6B	0.6063	0.2768	-0.2436	0.069*	
H6C	0.6419	0.3077	-0.3014	0.069*	
C7	0.62781 (7)	0.31108 (7)	-0.0377 (3)	0.0308 (6)	
C9	0.65864 (6)	0.28479 (7)	0.2463 (4)	0.0271 (5)	
C10	0.68264 (8)	0.25280 (8)	0.3421 (4)	0.0360 (6)	
H10A	0.6953	0.2648	0.4484	0.054*	
H10B	0.7040	0.2421	0.2631	0.054*	
H10C	0.6644	0.2305	0.3778	0.054*	

C8	0.64852 (7)	0.28190 (7)	0.0659 (3)	0.0303 (6)	
N2	0.66001 (8)	0.24416 (7)	-0.0254 (4)	0.0478 (6)	
O9	0.64699 (12)	0.21173 (8)	0.0305 (5)	0.0709 (11)	0.892 (7)
O10	0.68344 (13)	0.24671 (10)	-0.1541 (5)	0.0844 (13)	0.892 (7)
O9A	0.6323 (5)	0.2214 (6)	-0.064 (4)	0.047 (5)	0.108 (7)
O10A	0.6956 (5)	0.2340 (7)	-0.042 (5)	0.071 (6)	0.108 (7)
O5	0.67470 (5)	0.38351 (5)	0.1938 (2)	0.0318 (4)	
O6	0.62659 (5)	0.38295 (5)	0.4944 (3)	0.0309 (4)	
C11	0.72986 (8)	0.42775 (9)	0.1701 (4)	0.0436 (7)	
H11A	0.7548	0.4176	0.2274	0.065*	
H11B	0.7301	0.4577	0.1704	0.065*	
H11C	0.7287	0.4178	0.0468	0.065*	
C12	0.69317 (7)	0.41265 (7)	0.2705 (4)	0.0330 (6)	
C14	0.64901 (8)	0.41223 (7)	0.5448 (4)	0.0358 (7)	
C15	0.64112 (10)	0.42602 (9)	0.7323 (4)	0.0495 (8)	
H15A	0.6184	0.4456	0.7331	0.074*	
H15B	0.6657	0.4391	0.7801	0.074*	
H15C	0.6340	0.4023	0.8062	0.074*	
C13	0.67903 (11)	0.42915 (11)	0.4312 (5)	0.0372 (8)	0.848 (4)
N3	0.69807 (11)	0.46657 (10)	0.4996 (6)	0.0564 (9)	0.848 (4)
O11	0.73566 (11)	0.46798 (13)	0.5000 (8)	0.0929 (16)	0.848 (4)
O12	0.67605 (8)	0.49417 (7)	0.5453 (5)	0.0709 (11)	0.848 (4)
C13A	0.6847 (5)	0.4257 (5)	0.4490 (19)	0.0372 (8)	0.152 (4)
N3A	0.7157 (6)	0.4544 (5)	0.521 (3)	0.0564 (9)	0.152 (4)
O11A	0.7310 (5)	0.4489 (5)	0.665 (2)	0.073 (5)	0.152 (4)
O12A	0.7248 (7)	0.4836 (5)	0.433 (3)	0.073 (5)	0.152 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.02761 (17)	0.02064 (16)	0.02963 (16)	0.00184 (12)	-0.00204 (16)	-0.00131 (15)
O1	0.0348 (10)	0.0206 (8)	0.0479 (12)	0.0007 (7)	-0.0062 (8)	0.0027 (8)
O2	0.0318 (9)	0.0250 (9)	0.0416 (11)	-0.0002 (7)	-0.0010 (8)	-0.0023 (7)
C1	0.068 (2)	0.0282 (15)	0.056 (2)	0.0156 (14)	-0.0166 (17)	-0.0017 (13)
C2	0.0436 (15)	0.0251 (13)	0.0318 (15)	0.0094 (11)	-0.0087 (12)	-0.0073 (11)
C3	0.0296 (13)	0.0407 (14)	0.0325 (15)	0.0138 (11)	-0.0020 (13)	-0.0048 (13)
C4	0.0341 (14)	0.0344 (14)	0.0280 (16)	-0.0011 (11)	-0.0006 (11)	-0.0076 (11)
C5	0.0364 (15)	0.0536 (18)	0.0470 (18)	-0.0064 (14)	0.0034 (14)	-0.0052 (15)
N1	0.0492 (17)	0.0667 (16)	0.0451 (15)	0.0238 (13)	0.0066 (15)	0.0083 (15)
O7	0.087 (2)	0.0720 (16)	0.102 (2)	0.0464 (15)	0.0249 (16)	-0.0003 (15)
O8	0.0335 (12)	0.116 (2)	0.0715 (19)	0.0089 (14)	-0.0101 (12)	0.0049 (16)
O3	0.0397 (9)	0.0281 (9)	0.0290 (9)	0.0003 (7)	-0.0060 (8)	0.0002 (8)
O4	0.0359 (9)	0.0253 (9)	0.0263 (9)	0.0058 (7)	-0.0030 (8)	-0.0008 (7)
C6	0.0594 (18)	0.0432 (15)	0.0348 (16)	-0.0087 (13)	-0.0015 (15)	-0.0039 (15)
C7	0.0299 (13)	0.0299 (14)	0.0328 (15)	-0.0100 (10)	0.0023 (11)	-0.0017 (11)
C9	0.0229 (11)	0.0222 (11)	0.0360 (14)	-0.0025 (9)	0.0036 (11)	0.0003 (11)
C10	0.0351 (14)	0.0278 (13)	0.0451 (17)	0.0018 (11)	-0.0008 (13)	0.0036 (12)
C8	0.0342 (13)	0.0230 (13)	0.0338 (15)	-0.0024 (10)	0.0076 (11)	-0.0051 (10)
N2	0.0656 (17)	0.0331 (14)	0.0447 (16)	0.0041 (12)	0.0035 (14)	-0.0136 (12)

O9	0.114 (3)	0.0271 (14)	0.072 (2)	-0.0078 (15)	0.006 (2)	-0.0089 (15)
O10	0.121 (3)	0.063 (2)	0.069 (3)	0.0254 (19)	0.045 (2)	-0.0107 (17)
O9A	0.061 (10)	0.025 (9)	0.055 (11)	0.004 (8)	-0.020 (9)	-0.025 (8)
O10A	0.083 (11)	0.044 (10)	0.086 (13)	0.002 (9)	0.024 (10)	-0.038 (10)
O5	0.0301 (9)	0.0286 (9)	0.0366 (10)	-0.0006 (7)	0.0008 (8)	-0.0001 (8)
O6	0.0381 (9)	0.0234 (9)	0.0311 (11)	0.0006 (6)	-0.0008 (9)	-0.0033 (8)
C11	0.0358 (15)	0.0351 (15)	0.060 (2)	-0.0033 (12)	-0.0018 (14)	0.0147 (13)
C12	0.0280 (12)	0.0233 (12)	0.0477 (16)	0.0037 (9)	-0.0076 (13)	0.0081 (14)
C14	0.0441 (15)	0.0231 (13)	0.0402 (18)	0.0078 (11)	-0.0053 (12)	-0.0057 (12)
C15	0.068 (2)	0.0368 (15)	0.0434 (19)	0.0009 (13)	-0.0029 (16)	-0.0134 (14)
C13	0.0383 (18)	0.0222 (14)	0.0512 (19)	-0.0025 (13)	-0.0102 (15)	-0.0034 (13)
N3	0.0388 (19)	0.0424 (18)	0.088 (2)	-0.0136 (14)	-0.002 (2)	-0.0241 (19)
O11	0.0459 (19)	0.082 (3)	0.150 (5)	-0.0126 (18)	-0.016 (3)	-0.050 (3)
O12	0.0668 (16)	0.0290 (14)	0.117 (3)	-0.0038 (13)	0.0069 (18)	-0.0255 (16)
C13A	0.0383 (18)	0.0222 (14)	0.0512 (19)	-0.0025 (13)	-0.0102 (15)	-0.0034 (13)
N3A	0.0388 (19)	0.0424 (18)	0.088 (2)	-0.0136 (14)	-0.002 (2)	-0.0241 (19)
O11A	0.051 (8)	0.081 (9)	0.087 (10)	-0.017 (7)	-0.035 (8)	-0.021 (8)
O12A	0.068 (10)	0.064 (9)	0.086 (10)	-0.043 (8)	-0.006 (8)	-0.017 (8)

Geometric parameters (Å, °)

Co1—O1	1.8635 (16)	C9—C10	1.493 (3)
Co1—O5	1.8686 (17)	C10—H10A	0.9800
Co1—O6	1.869 (2)	C10—H10B	0.9800
Co1—O4	1.8694 (16)	C10—H10C	0.9800
Co1—O2	1.8722 (17)	C8—N2	1.461 (3)
Co1—O3	1.8721 (19)	N2—O9A	1.208 (14)
O1—C2	1.261 (3)	N2—O10A	1.216 (15)
O2—C4	1.263 (3)	N2—O9	1.217 (4)
C1—C2	1.512 (3)	N2—O10	1.235 (4)
C1—H1A	0.9800	O5—C12	1.267 (3)
C1—H1B	0.9800	O6—C14	1.264 (3)
C1—H1C	0.9800	C11—C12	1.501 (4)
C2—C3	1.403 (4)	C11—H11A	0.9800
C3—C4	1.412 (4)	C11—H11B	0.9800
C3—N1	1.465 (3)	C11—H11C	0.9800
C4—C5	1.495 (4)	C12—C13	1.399 (5)
C5—H5A	0.9800	C12—C13A	1.432 (13)
C5—H5B	0.9800	C14—C13	1.412 (5)
C5—H5C	0.9800	C14—C13A	1.439 (13)
N1—O8	1.224 (4)	C14—C15	1.499 (4)
N1—O7	1.224 (3)	C15—H15A	0.9800
O3—C7	1.270 (3)	C15—H15B	0.9800
O4—C9	1.264 (3)	C15—H15C	0.9800
C6—C7	1.485 (4)	C13—N3	1.466 (4)
C6—H6A	0.9800	N3—O12	1.204 (4)
C6—H6B	0.9800	N3—O11	1.230 (5)
C6—H6C	0.9800	C13A—N3A	1.482 (16)
C7—C8	1.405 (4)	N3A—O12A	1.201 (18)
C9—C8	1.395 (4)	N3A—O11A	1.202 (18)

O1—Co1—O5	87.03 (7)	O4—C9—C10	114.4 (2)
O1—Co1—O6	87.82 (8)	C8—C9—C10	122.9 (2)
O5—Co1—O6	94.70 (7)	C9—C10—H10A	109.5
O1—Co1—O4	174.00 (8)	C9—C10—H10B	109.5
O5—Co1—O4	88.92 (7)	H10A—C10—H10B	109.5
O6—Co1—O4	88.11 (7)	C9—C10—H10C	109.5
O1—Co1—O2	95.40 (7)	H10A—C10—H10C	109.5
O5—Co1—O2	176.10 (8)	H10B—C10—H10C	109.5
O6—Co1—O2	88.46 (8)	C9—C8—C7	127.2 (2)
O4—Co1—O2	88.89 (7)	C9—C8—N2	116.8 (2)
O1—Co1—O3	88.84 (8)	C7—C8—N2	116.1 (2)
O5—Co1—O3	88.69 (8)	O9A—N2—O10A	121.6 (13)
O6—Co1—O3	175.11 (7)	O9—N2—O10	123.0 (3)
O4—Co1—O3	95.49 (7)	O9A—N2—C8	116.2 (9)
O2—Co1—O3	88.30 (8)	O10A—N2—C8	121.6 (10)
C2—O1—Co1	126.46 (16)	O9—N2—C8	119.0 (3)
C4—O2—Co1	126.32 (16)	O10—N2—C8	117.9 (3)
C2—C1—H1A	109.5	C12—O5—Co1	124.92 (17)
C2—C1—H1B	109.5	C14—O6—Co1	124.96 (18)
H1A—C1—H1B	109.5	C12—C11—H11A	109.5
C2—C1—H1C	109.5	C12—C11—H11B	109.5
H1A—C1—H1C	109.5	H11A—C11—H11B	109.5
H1B—C1—H1C	109.5	C12—C11—H11C	109.5
O1—C2—C3	122.6 (2)	H11A—C11—H11C	109.5
O1—C2—C1	112.4 (2)	H11B—C11—H11C	109.5
C3—C2—C1	125.0 (2)	O5—C12—C13	121.6 (3)
C2—C3—C4	126.1 (2)	O5—C12—C13A	123.8 (6)
C2—C3—N1	118.1 (2)	O5—C12—C11	113.6 (3)
C4—C3—N1	115.8 (2)	C13—C12—C11	124.7 (3)
O2—C4—C3	122.5 (2)	C13A—C12—C11	121.7 (6)
O2—C4—C5	113.8 (2)	O6—C14—C13	121.3 (3)
C3—C4—C5	123.6 (2)	O6—C14—C13A	123.6 (7)
C4—C5—H5A	109.5	O6—C14—C15	114.1 (3)
C4—C5—H5B	109.5	C13—C14—C15	124.6 (3)
H5A—C5—H5B	109.5	C13A—C14—C15	121.0 (7)
C4—C5—H5C	109.5	C14—C15—H15A	109.5
H5A—C5—H5C	109.5	C14—C15—H15B	109.5
H5B—C5—H5C	109.5	H15A—C15—H15B	109.5
O8—N1—O7	124.2 (3)	C14—C15—H15C	109.5
O8—N1—C3	119.0 (3)	H15A—C15—H15C	109.5
O7—N1—C3	116.8 (3)	H15B—C15—H15C	109.5
C7—O3—Co1	126.27 (17)	C12—C13—C14	126.7 (3)
C9—O4—Co1	126.28 (17)	C12—C13—N3	118.9 (3)
C7—C6—H6A	109.5	C14—C13—N3	114.3 (3)
C7—C6—H6B	109.5	O12—N3—O11	124.7 (3)
H6A—C6—H6B	109.5	O12—N3—C13	118.1 (3)
C7—C6—H6C	109.5	O11—N3—C13	117.2 (4)
H6A—C6—H6C	109.5	C12—C13A—C14	122.1 (11)

H6B—C6—H6C	109.5	C12—C13A—N3A	113.4 (11)
O3—C7—C8	122.1 (2)	C14—C13A—N3A	124.4 (12)
O3—C7—C6	114.9 (2)	O12A—N3A—O11A	120.5 (19)
C8—C7—C6	123.0 (2)	O12A—N3A—C13A	118.3 (17)
O4—C9—C8	122.6 (2)	O11A—N3A—C13A	121.2 (15)
O5—Co1—O1—C2	-177.2 (2)	C7—C8—N2—O9	-120.7 (4)
O6—Co1—O1—C2	-82.3 (2)	C9—C8—N2—O10	-120.0 (4)
O2—Co1—O1—C2	5.9 (2)	C7—C8—N2—O10	61.0 (4)
O3—Co1—O1—C2	94.1 (2)	O1—Co1—O5—C12	65.7 (2)
O1—Co1—O2—C4	1.0 (2)	O6—Co1—O5—C12	-21.9 (2)
O6—Co1—O2—C4	88.7 (2)	O4—Co1—O5—C12	-109.90 (19)
O4—Co1—O2—C4	176.8 (2)	O3—Co1—O5—C12	154.58 (19)
O3—Co1—O2—C4	-87.6 (2)	O1—Co1—O6—C14	-64.51 (19)
Co1—O1—C2—C3	-10.3 (4)	O5—Co1—O6—C14	22.32 (19)
Co1—O1—C2—C1	170.89 (18)	O4—Co1—O6—C14	111.09 (19)
O1—C2—C3—C4	7.7 (4)	O2—Co1—O6—C14	-159.97 (19)
C1—C2—C3—C4	-173.7 (3)	Co1—O5—C12—C13	8.8 (4)
O1—C2—C3—N1	-174.7 (3)	Co1—O5—C12—C13A	20.7 (10)
C1—C2—C3—N1	3.9 (4)	Co1—O5—C12—C11	-169.52 (16)
Co1—O2—C4—C3	-3.4 (4)	Co1—O6—C14—C13	-9.6 (4)
Co1—O2—C4—C5	178.68 (17)	Co1—O6—C14—C13A	-21.4 (10)
C2—C3—C4—O2	-0.3 (4)	Co1—O6—C14—C15	171.42 (18)
N1—C3—C4—O2	-178.0 (3)	O5—C12—C13—C14	12.8 (5)
C2—C3—C4—C5	177.4 (3)	C13A—C12—C13—C14	-93 (4)
N1—C3—C4—C5	-0.3 (4)	C11—C12—C13—C14	-169.1 (3)
C2—C3—N1—O8	133.5 (3)	O5—C12—C13—N3	-171.0 (3)
C4—C3—N1—O8	-48.7 (4)	C13A—C12—C13—N3	83 (4)
C2—C3—N1—O7	-46.8 (4)	C11—C12—C13—N3	7.0 (5)
C4—C3—N1—O7	131.1 (3)	O6—C14—C13—C12	-12.4 (5)
O1—Co1—O3—C7	173.7 (2)	C13A—C14—C13—C12	94 (4)
O5—Co1—O3—C7	86.68 (19)	C15—C14—C13—C12	166.5 (3)
O4—Co1—O3—C7	-2.11 (19)	O6—C14—C13—N3	171.4 (3)
O2—Co1—O3—C7	-90.83 (19)	C13A—C14—C13—N3	-83 (4)
O5—Co1—O4—C9	-84.85 (18)	C15—C14—C13—N3	-9.8 (5)
O6—Co1—O4—C9	-179.59 (18)	C12—C13—N3—O12	130.7 (4)
O2—Co1—O4—C9	91.91 (19)	C14—C13—N3—O12	-52.7 (6)
O3—Co1—O4—C9	3.73 (19)	C12—C13—N3—O11	-46.7 (7)
Co1—O3—C7—C8	0.4 (3)	C14—C13—N3—O11	129.9 (5)
Co1—O3—C7—C6	179.31 (17)	O5—C12—C13A—C14	-12.7 (19)
Co1—O4—C9—C8	-3.6 (3)	C13—C12—C13A—C14	68 (4)
Co1—O4—C9—C10	174.29 (15)	C11—C12—C13A—C14	178.3 (9)
O4—C9—C8—C7	0.8 (4)	O5—C12—C13A—N3A	168.2 (9)
C10—C9—C8—C7	-176.8 (2)	C13—C12—C13A—N3A	-111 (5)
O4—C9—C8—N2	-178.0 (2)	C11—C12—C13A—N3A	-0.8 (15)
C10—C9—C8—N2	4.3 (3)	O6—C14—C13A—C12	13.1 (19)
O3—C7—C8—C9	0.9 (4)	C13—C14—C13A—C12	-67 (4)
C6—C7—C8—C9	-178.0 (2)	C15—C14—C13A—C12	179.4 (10)
O3—C7—C8—N2	179.7 (2)	O6—C14—C13A—N3A	-167.9 (11)

C6—C7—C8—N2	0.9 (4)	C13—C14—C13A—N3A	112 (5)
C9—C8—N2—O9A	108.5 (17)	C15—C14—C13A—N3A	-1.6 (18)
C7—C8—N2—O9A	-70.5 (17)	C12—C13A—N3A—O12A	51 (2)
C9—C8—N2—O10A	-63 (2)	C14—C13A—N3A—O12A	-128 (2)
C7—C8—N2—O10A	118 (2)	C12—C13A—N3A—O11A	-130.2 (19)
C9—C8—N2—O9	58.3 (4)	C14—C13A—N3A—O11A	51 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C1—H1B...O7 ⁱ	0.98	2.33	3.087 (4)	134
C11—H11B...O12 ⁱⁱ	0.98	2.55	3.240 (4)	128
C10—H10C...O5 ⁱⁱⁱ	0.98	2.46	3.433 (3)	176
C15—H15C...O3 ^{iv}	0.98	2.57	3.542 (4)	174

Symmetry codes: (i) $-x+1, y, z-1/2$; (ii) $x, -y+1, z-1/2$; (iii) $-y+1, x-1/2, z+1/4$; (iv) $x, y, z+1$.